

AD-A174 871

APPEARANCE OF INGESTED H SUB 2 (18)O IN PLASMA AND  
SWEAT DURING EXERCISE- (U) ARMY RESEARCH INST OF  
ENVIRONMENTAL MEDICINE NATICK MA L E ARMSTRONG ET AL  
24 NOV 86 USARIEM-M-11-87


1/1

UNCLASSIFIED

F/G 6/19

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS 1963-A

1

1

APPEARANCE OF INGESTED  $H_2^{18}O$  IN PLASMA AND SWEAT  
DURING EXERCISE-HEAT EXPOSURE

Authors: Lawrence E. Armstrong, Ph.D.

Roger V. Hubbard, Ph.D.

Steven H. Zeisel, M.D.

Morteza Janghorbani, Ph.D.

Institutions: USARIEM, Heat Research Division, Natick, MA 01760-5007.

and

Boston University School of Medicine, Boston, MA 02118.

Correspondence: Lawrence E. Armstrong

Research Physiologist

Heat Research Division

U. S. Army Research Institute of Environmental Medicine

Natick, MA 01760-5007.

(617) 651-4875

Running Head: Stable Isotopes in Plasma and Sweat

**DISTRIBUTION STATEMENT A**

Approved for public release;  
Distribution Unlimited

(1 table, 2 figures)

DTIC  
ELECTE  
DEC 5 1986  
S B

86 12 05 043

AD-A174 871

DTIC FILE COPY

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No 0704-0188  
Exp Date Jun 30, 1986

1a. REPORT SECURITY CLASSIFICATION <b>UNCLAS</b>			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT DISTRIBUTION A: Approved for public release; distribution is unlimited.		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) <b>M11-87</b>			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Heat Research Division		6b. OFFICE SYMBOL (If applicable) SGRD-UE-HR	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) US Army Research Institute of Environmental Medicine Natick, MA 01760-5007			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO. 129
11. TITLE (Include Security Classification) Appearance of ingested $H_2^{18}O$ in plasma and sweat during exercise-heat exposure					
12. PERSONAL AUTHOR(S) L.E. Armstrong, R.W. Hubbard, S.H. Zeisel, M. Janghorbani					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) 86 11 24	
15. PAGE COUNT 16					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) In an effort to study human water transport and eccrine sweat gland function, this investigation measured the rate of appearance of $H_2^{18}O$ in plasma and sweat. Four healthy males were exposed for 6 h to a hot, wet environment (37.1°C, db, 31.3°C wb) and to intermittent cycle ergometer protocols. Baseline plasma (antecubital vein) and scraped sweat samples were collected (-30 to 0 min) prior to administration of 100 g of labelled water (84.7% $^{18}O$ , 15.3% $^{16}O$ ) via nasogastric tube at 0 min. Samples were analyzed using isotopic ratio mass spectrometry. The isotopic enrichment of baseline sweat samples was slightly greater than that of plasma samples. Peak enrichment ( $^{18}O/^{16}O$ ratio) in plasma (range: $2.4908 - 2.7206 \times 10^{-3}$ ) occurred at 21 - 28 min postdose and at 21 - 45 min postdose in sweat (range: $2.3089 - 2.6666 \times 10^{-3}$ ). The $^{18}O/^{16}O$ ratio in plasma and sweat declined rapidly, then declined slowly for the remaining heat exposure. The appearance of $H_2^{18}O$ in sweat reflected that of plasma; neither curve was significantly altered by exercise intensity, duration, or frequency. To our knowledge, this is the first stable isotope data to verify that ingested fluid is rapidly assimilated and becomes available for evaporative cooling during work in the heat.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION		
22a. NAME OF RESPONSIBLE INDIVIDUAL			22b. TELEPHONE (Include Area Code)		22c. OFFICE SYMBOL

APPEARANCE OF INGESTED  $H_2^{18}O$  IN PLASMA AND SWEAT  
DURING EXERCISE-HEAT EXPOSURE



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
<b>PER FORM 50</b>	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
<b>A-1</b>	

(1 table, 2 figures)

Abstract

Oxygen 16

Oxygen 18

In an effort to study human water transport and eccrine sweat gland function, this investigation measured the rate of appearance of  $H_2^{18}O$  in plasma and sweat. Four healthy males were exposed for 6 h to a hot, wet environment ( $37.1^\circ C$  db,  $31.3^\circ C$  wb) and to intermittent cycle ergometer protocols. Baseline plasma (antecubital vein) and scraped sweat samples were collected (-30 to 0 min) prior to administration of 100 g of labelled water ( $84.7\% \text{ }^{16}O$ ,  $15.3\% \text{ }^{18}O$ ) via nasogastric tube at 0 min. Samples were analyzed using isotopic ratio mass spectrometry. The isotopic enrichment of baseline sweat samples was slightly greater than that of plasma samples. Peak enrichment ( $^{18}O/^{16}O$  ratio) in plasma (range:  $2.4908 - 2.7206 \times 10^{-3}$ ) occurred at 21 - 28 min postdose and at 21 - 45 min postdose in sweat (range:  $2.3089 - 2.6666 \times 10^{-3}$ ). The  $^{18}O/^{16}O$  ratio in plasma and sweat declined rapidly, then declined slowly for the remaining heat exposure. The appearance of  $H_2^{18}O$  in sweat reflected that of plasma; neither curve was significantly altered by exercise intensity, duration, or frequency. To our knowledge, this is the first stable isotope data to verify that ingested fluid is rapidly assimilated and becomes available for evaporative cooling during work in the heat.

KEY WORDS: stable isotope, blood plasma, sweat,  $H_2^{18}O$ , fluid transport, sweat gland

## Introduction

Eccrine sweat glands have been studied for a variety of reasons (6,9,12,16), the most important of which is to measure the effects of sweat secretion on whole body fluid/electrolyte balance and thermal balance. Early studies focused on the effects of water intake on sweat rate; Lee and Mulder (8), for example, were the first to report that bursts of sweating could be observed approximately 3 minutes after water consumption. Similar bursts of sweating have been described by Kuno (6), Senay and Christensen (15), Banerjee and Bullard (3), and Lechner and Hertig (7). However, evidence is still lacking to determine whether these bursts represent a reflex response to fluid intake, or are a manifestation of altered pressure/volume relationships or osmolarity within the circulatory system. Researchers routinely have assumed that a sweating onset of 2-3 minutes is better explained by a reflex action than by an increase of either the vascular or interstitial space (3,6,7,8). To date, no data exist regarding the time course of the appearance of an ingested fluid in eccrine sweat.

The exercising adult can benefit from such data, as well. The loss of water by eccrine sweating is often greater than the intake of water by drinking (10). This is especially true when exercise intensities exceed 70%  $\text{VO}_2$  max; at that point, gastric emptying appears to become an important limiting factor (4). One recent report, for example, described the unavoidable body weight deficits which a world class distance runner experienced during a hot, humid marathon competition (2), in spite of his efforts to prehydrate and to drink often. The common advice given to athletes and fitness enthusiasts is to drink during training and competition; yet, no data exist to indicate exactly how rapidly ingested water appears in eccrine

sweat or whether this advice is necessary during brief, intermittent exercise bouts. The advice to drink during exercise may offer protection against dehydration, but may offer little additional benefit from evaporative cooling. An improved understanding of the availability of ingested water for use by eccrine sweat glands may even modify current heat injury prevention doctrine.

Although eccrine sweat gland research is a relatively unexplored area of physiological research when compared to the recent progress being made in the study of other transport epithelia (12), the two aforementioned problems have not been investigated because the appropriate technology was not available and because radioactive tracers are undesirable in human studies. With the advent of stable isotope tracer methodology, stable isotopes such as  $^2\text{H}$  or  $^{18}\text{O}$  have proven to be suitable for measurements in a variety of biological fluids (14, Janghorbani unpublished observations.) Despite obvious value, stable isotope tracer methods have seldom been used in relation to the dynamics of body water during exercise, or in evaluating the function of eccrine sweat glands. Therefore, the purpose of the present investigation was to measure the rate of appearance of ingested water (using the stable isotope  $\text{H}_2^{18}\text{O}$ ) in plasma and eccrine sweat, during intermittent exercise in the heat. Four males were exposed to an ambient temperature of  $37.1^\circ\text{C}$  for 6 hours. Following the ingestion of an  $\text{H}_2^{18}\text{O}$  labelled drink, blood and sweat samples were collected periodically. By measuring the isotope ratio of  $^{18}\text{O}$  to  $^{16}\text{O}$ , the rate of entry of the labelled drink was analyzed in eccrine sweat and in plasma.



### Material & Methods

The subjects of this investigation were four healthy, unacclimatized males who received a thorough explanation of techniques, potential risks and benefits, prior to giving their voluntary written consent to participate. Physical characteristics of subjects A, B, C, and D, respectively, were as follows: age - 37, 43, 47, 28 yr; height - 166, 178, 168, 184 cm; mass - 66.757, 58.822, 88.049, 94.250 Kg; surface area - 1.84, 1.67, 1.96, 2.14 m<sup>2</sup>. Prior to arriving at the laboratory at 0800h, subjects had eaten no food for at least 2 hours. Electrocardiograph leads, a rectal thermister (8cm beyond the anal sphincter), and a forearm intravenous catheter were fitted after subjects entered the environmental chamber (37.1  $\pm$  0.2°C dry bulb, 31.3  $\pm$  0.2°C wet bulb). Heart rate, rectal temperature, body weight ( $\pm$  10g), and blood pressure (sphygmomanometer) were recorded at regular intervals during each six hour exercise-heat exposure. Sweat rate was calculated by using body weight differences, corrected ( $\pm$  10g) for water and food intake as well as urine output. Exercise was utilized primarily as a stimulus for sweat production. All subjects exercised on a bicycle ergometer (Monark, model 868, Stockholm, Sweden) at work rates which were individually selected.

Baseline blood and sweat samples were obtained prior to the ingestion of 100g H<sub>2</sub><sup>18</sup>O (84.7% <sup>16</sup>O and 15.3% <sup>18</sup>O, Mound Laboratories, Miamisburg, OH). This bolus of labelled water was delivered via nasogastric tube and was rinsed with 100g distilled water. Water was consumed ad libitum and a light lunch and beverages were ingested between 3.5-4.5 hours postdosing. During the initial hour, blood and sweat samples were obtained at 10-15 min intervals, then at 60 min intervals for the remaining five hours. Venous blood was sampled from an antecubital vein in heparinized tubes, centrifuged, and the

plasma fraction was retained for isotopic analysis. Sweat (2-5ml) was scraped from the forehead and neck into clean polyethylene tubes. No attempt was made to collect uncontaminated samples because this sweat sampling technique was being evaluated for possible use in later field and laboratory studies.

The  $^{18}\text{O}/^{16}\text{O}$  ratios in blood and sweat samples were analyzed using isotope ratio mass spectrometry (Isogas 903, Vacuum Generators, Manchester, U.K.). The rise in the  $^{18}\text{O}/^{16}\text{O}$  ratio for either blood or sweat indicated the rate of entry of the  $^{18}\text{O}$  enriched water for that fluid. Values were expressed as the g-atom ratio of  $^{18}\text{O}/^{16}\text{O}$  by the use of a set of water standards spiked with varying but known amounts of  $\text{H}_2^{18}\text{O}$ . The measurement precision for  $^{18}\text{O}/^{16}\text{O}$  ratios in deionized water and various physiological fluids was evaluated at various levels of isotopic enrichment. The standard deviation for seven triplicate sets of these fluids was within the range 0.005 - 0.06%.

### Results

#### FIGURE 1

Figure 1 depicts the duration of exercise and the hourly work output for each subject. Because bicycle ergometry was utilized to promote adequate sweat production, and because catheter patency, hourly sample collection and symptoms of heat illness were of primary importance, little effort was made to maintain a regular exercise schedule. For example, subjects A and B performed 43-46 min of exercise during the initial 60 min postdosing, while subjects C and D did no work on the ergometer. Subjects A and B also exercised during every hourly segment (exception: hour 2 for subject B), whereas subjects C and D exercised only during hours 2, 3 and 4. Steady-state heart rates during exercise for subjects A, B, C and D ranged from: 102-120, 114-126, 88-100, 84-120 beats  $\text{min}^{-1}$ . Sweat rates were stable after the initial hour, and averaged  $223 \pm 25$ ,  $105 \pm 35$ ,  $223 \pm 51$ , and  $260 \pm 71 \text{ ml} \cdot \text{m}^{-2} \cdot \text{h}^{-1}$  (subjects A, B,

C and D respectively). The mean change in rectal temperature was  $0.2 \pm 0.2^{\circ}\text{C}$  during these six hour trials.

TABLE 1 Table 1 describes baseline isotope ratios, prior to ingestion of the 100ml dose of  $^{18}\text{O}$  labelled water. Sweat samples exhibited higher  $^{18}\text{O}/^{16}\text{O}$  ratios than did corresponding plasma samples.

FIGURE 2 Figure 2 compares rate of entry of  $^{18}\text{O}$  labelled water into plasma and sweat. The isotope ratio in plasma rose sharply, with peak appearance occurring at 21-28 min. A decline was observed in plasma until 60-150 min, depending on the subject. Thereafter, the plasma isotope ratio declined gradually for the remainder of the six hour observation period. Although the postdose isotopic enrichment was always higher in plasma, the rate of appearance of  $^{18}\text{O}$  in sweat essentially paralleled its appearance in the plasma of all subjects. The time of  $^{18}\text{O}$  peak appearance in sweat ranged from 18 min (subject D) to 45 min (subject C). Evidence of exogenous contamination (Fig. 2) appeared randomly at points in the sweat curves of subject C (275 min) and subject D (43 min).

#### Discussion

The isotopic enrichment of baseline sweat samples was greater than that of baseline plasma samples (Table 1). The exact reason for this observation in all four subjects is not evident, but these values are consistent with greater evaporation of the lighter isotope ( $^{16}\text{O}$ ). An alternative explanation involves preferential secretion of  $^{18}\text{O}$  by eccrine sweat glands. All plasma samples taken after the ingestion of labelled water (fig. 2) exhibited higher isotopic ratios than sweat. The rapid rise and subsequent fall of the plasma and sweat isotope ratios (fig. 2) represented first the rate of absorption and subsequently the equilibration of  $\text{H}_2^{18}\text{O}$  in the vascular, interstitial and

intracellular compartments. Once the plasma and sweat curves had plateaued (approximately 60-120 min), a gradual decline was observed in the  $^{18}\text{O}/^{16}\text{O}$  ratio (Fig. 2) for the remainder of the six hours. This gradual decline was probably due to four factors: (a) dilution with ad libitum liquid intake, (b) mixing of total body water with those compartments which are not readily accessible (e.g. synovial fluid, cerebrospinal fluid), (c) true expansion of body water from metabolically generated water, and (d) other anomalies such as isotope exchange between body water and the bicarbonate pool or possible exchange with nonaqueous oxygen.

The production of sweat in bursts shortly after water consumption has been observed by several investigators (3,6,7,8) but data are lacking to indicate whether these bursts represent a reflex response or an altered pressure/volume relationship within the circulatory system. The present investigation demonstrates that these theories may be examined using the stable isotope techniques described above. Oral administration of  $\text{H}_2^{18}\text{O}$  would not be appropriate for short term measurements (e.g. 3 min postdose), however. Injection or infusion of the labelled water would be necessary.

Loss of water during exercise in the heat often exceeds the water replaced by drinking (2,10). The advice to consume water during exercise may offer protection against dehydration, but the time course of water availability for use by eccrine sweat glands was not known prior to this investigation. Figure 2 indicates that the  $^{18}\text{O}$  labelled water appeared in sweat shortly after ingestion (9-18 min) and was maximally present in sweat at 21-45 minutes, depending on the subject. Because sweat rate may decline with severe dehydration (5,11,13), ingested water can clearly aid the exercising adult by maintaining skin wettedness during prolonged exercise. This is especially

significant considering the fact that evaporation of sweat may account for 85-96% of total heat dissipation during strenuous exercise in dry heat (1).

Comparison of Figure 1 with Figure 2 indicates that the appearance of  $\text{H}_2^{18}\text{O}$  in sweat was not influenced by the frequency, duration, or intensity of exercise, but apparently was solely a function of the rate of entry into the vascular space. The  $^{18}\text{O}$  appearance curves of all subjects were strikingly similar, in spite of varying exercise regimes (Fig. 1) and ad libitum water intake.

Differences in isotopic ratios among various body fluids have been observed previously (14), and sweat is among the least preferred physiological matrices because of its potential for exogenous contamination (Janghorbani, unpublished observations). By sampling from enclosed skin surfaces, it should be possible to eliminate contamination. However, the data herein have successfully described the time course of fluid movements between the gastrointestinal tract, the vascular compartment and the interstitial space, and has clarified the physiological function of eccrine sweat glands. This investigation also has demonstrated that stable isotope methods can be valuable in measuring fluid dynamics during future heat acclimation, physical training, disease, and nutrition studies.

## References

1. Adams, W.C., R.H. Fox, A.J. Fry, and I.C. MacDonald. Thermoregulation during marathon running in cool, moderate and hot environments. J. Appl. Physiol. 38 (6):1030-1037, 1975.
2. Armstrong, L.E., R.W. Hubbard, B.H. Jones, J.T. Daniels. Preparing Alberto Salazar for the heat of the 1984 Olympic marathon. Physician and Sportsmedicine 14(3):73-81, 1986.
3. Banerjee, M.R., and R.W. Bullard. Transient changes in sweating activity following water ingestion. Biometeorology 4(2):15, 1969.
4. Costill, D.L., and B. Saltin. Factors limiting gastric emptying during rest and exercise. J. Appl. Physiol. 37:679-683, 1974.
5. Fortney, S.M., E.R. Nadel, C.B. Wenger, and J.R. Bove. Effect of acute alterations of blood volume on circulatory performance in humans. J. Appl. Physiol. 50:292-298, 1981.
6. Kuno, Y. Human Perspiration. Springfield, IL: Charles C. Thomas Publishers, 1956, p. 264-270.
7. Lechner, P.L., and B.A. Hertig. Effect of water intake on sweat rate. Fed. Proc. 28:528, 1969.
8. Lee, D.H.K., and A.G. Mulder. Some immediate effects of reduced cooling powers upon the water balance and related effects in the human subject. J. Physiol. (London) 84:410-432, 1935.
9. Robinson, S., and A.H. Robinson. Chemical composition of sweat. Physiol. Rev. 34:202-220, 1954.
10. Rothstein, A., E.F. Adolph, and J.H. Wills. Voluntary dehydration. In: Physiology of Man in the Desert, edited by E.F. Adolph and associates. New York: Interscience, 1947, p. 254-270.


11. Sargent, F. II, W.K. Brown, A.T. Pessa, J. Waligora, and K.P. Weinman. Observations on dehydration and eccrine sweating. In: Environmental Physiology and Psychology in Arid Conditions. Leige, Belgium: UNESCO, 1964, p. 115-119.
12. Sato, K. The physiology, pharmacology, and biochemistry of the eccrine sweat gland. Rev. Physiol. Biochem. Pharmacol. 79:51-131, 1977.
13. Sawka, M.N., A.J. Young, R.P. Francesconi, S.R. Muza, and K.B. Pandolf. Thermoregulatory and blood responses during exercise at graded hypohydration levels. J. Appl. Physiol. 59(5):1394-1401, 1985.
14. Schoeller, D.A., E. van Santen, D.W. Peterson, W. Dietz, J. Jaspán, and P.D. Klein. Total body water measurement in humans with  $^{18}\text{O}$  and  $^2\text{H}$  labelled water. Am. J. Clin. Nutr. 33:2686-2693, 1980.
15. Senay, L.C., and M.L. Christensen. Cardiovascular and sweating responses to water ingestion during dehydration. J. Appl. Physiol. 20:975-979, 1965.
16. Weiner, J.S., and K. Hellman. The sweat glands. Biol. Rev. 35:141-187, 1960.

Figure Titles

Figure 1 - Duration, hourly work output and total work performed by subjects A - D.

Figure 2 - Appearance of  $\text{H}_2^{18}\text{O}$  in plasma and sweat of subjects A - D.





\* brief syncope

● PLASMA  
— SWEAT







Table 1 - Observed isotope ratios ( $^{18}\text{O}/^{16}\text{O}$ ) in predose (0 min) samples of plasma and sweat.

<u>Subject</u>	<u>Isotope ratio</u>	
	<u>Plasma</u>	<u>Sweat</u>
A	$2.0506 \times 10^{-3}$	$2.0677 \times 10^{-3}$
B	$2.0522 \times 10^{-3}$	$2.0637 \times 10^{-3}$
C	$2.0493 \times 10^{-3}$	$2.0614 \times 10^{-3}$
D	$2.0503 \times 10^{-3}$	$2.0586 \times 10^{-3}$

END

1-87

DTIC